

## Effects of gravitational unloading on sensory-motor system

I. B. Kozlovskaya; Moskau

In the multicomponent system of motor control microgravity results in several effects which can alter dramatically characteristics of acitivity of various specific mechanisms and the systems as a whole. Quantitative studies of changes caused by short and long-term exposures to real (spaceflight, SF) and simulated (dry immersion, DI, and antiorthostatic bedrest, ABR) microgravity in different parts and mechanisms of the motor systems demonstrated deep alterations in 1) acivities of different sensory inputs; 2) morphological and physiological characteristics of muscles; 3) functioning of almost all motor control mechanisms under study. The depth of alterations as well as combinations of changes after exposures of different durations varied considerable pointing out to different nature of disturbances at varying stages of exposure to microgravity.

The main features of short-term effects revealed during and after 7-day DI, 30 day ABR and 7-13 day SF were: 1) different inputs and reflex mechanisms hypersensitivity which, seemed, to have non-specific character and was demonstrated clearly by changes in parameters of a) H- and T-spinal reflexes of musculus triceps surae; b) vestibulo-ocular reactions both voluntary and evoked; c) motor units activity and 2) sharp drop of extensor muscle tone causing in 1st turn a considerable decrease of a) muscle strength (up to 30%), b) movements precision and c) orthostatic tolerance. After exposure of longer durations (data of 120 and 370 days ABR and long-term SF) the reflex hypersensitivity clearly diminished being suppressed by active inhibitory processes. Muscle atrophic processes revealed by progressive diminishing of extensor muscle strength (sometimes flexor muscles too), a decrease of M-response amplitude and deep changes of motor unit rhythmic activity as well as significant alterations of central motor control mechanisms activity due partially of adaptations processes seemed to contribute mainly to the development of movements disorders at this stage.

## Untersuchungen in der simulierten und echten Schwerelosigkeit, Anwendungsmöglichkeiten in Diagnostik und Therapie bei neurologischen Erkrankungen

F. Gerstenbrand, M. Marosi; Innsbruck

Beim Start einer Trägerrakete zum Raumflug und kurz nach Eintreten in die Schwerelosigkeit kann sich bei einzelnen Astronauten und Kosmonauten ein „Space-adaption-syndrome“ mit den im Vordergrund stehenden Symptomen der „motion sickness“ einstellen, das sich in vegetativen Symptomen, Bewegungsstörungen bis zur Ataxie optomotorischen Ausfällen, kognitiver Leistungsminderung etc. zeigt. Nach längerem Aufenthalt in der Schwerelosigkeit entsteht das Kosmonautensyndrom mit spinaler Ataxie, zerebellärer Ataxie, primärer Muskelatrophie, Polyneuropathie, verminderter Vigilanz, Störungen des Körperschemas und der höheren sowie der höchsten kortikalen Funktionen, aber auch vegetativen Veränderungen und Osteoporose etc.

Der Zustand der Schwerelosigkeit kann im Schwerefeld der Erde mittels Parabolflügen dem Bed-Rest Modell sowie dem Dry-Water-Immersion-Modell partiell simuliert werden. Die in der simulierten Schwerelosigkeit auftretenden Störungen entsprechen weitgehend dem Kosmonautensyndrom und lassen sich klinisch als Bed-Rest-Syndrom zusammenfassen. Als Krankheitsbild tritt das Bed-Rest-Syndrom bei langer Bettlägerigkeit wie bei Komazuständen verschiedener Ätiologie, nach Wirbel- oder Beckenfraktur, kardialer Insuffizienz, aber auch bei älteren Menschen mit verminderter Bewegungsintention obligatorisch auf.

Durch die Untersuchungen in der echten und in der simulierten Schwerelosigkeit konnten neue Erkenntnisse über die menschliche Motorik, sowie über die kognitiven Funktionen des Menschen, vor allem über seine psychomotorischen Reaktionen, die Kontrolle von Emotionen, ferner über die

Sprachfunktionen, die Raumorientierungsfähigkeit u. a. gewonnen werden.

Neue Erkenntnisse ergaben sich über den Muskelstoffwechsel. Der Forschungsbereich des Flüssigkeitstransportes in der Schwerelosigkeit hat ebenfalls interessante Erkenntnisse erbracht und wichtige Voraussetzungen für den Aufenthalt des Menschen im Weltraum geschaffen. Die Erfahrungen aus der Forschung in der echten und der simulierten Schwerelosigkeit sind in die Routine medizin, und zwar in die Prävention, die Therapie und die Rehabilitation sowohl von neurologischen als auch von anderen Erkrankungen zu übertragen. So konnten neue Verfahren für die Behandlung von Hirnverletzten, Schlaganfallpatienten etc. entwickelt werden. Mit Hilfe der Untersuchungsmethoden der simulierten Schwerelosigkeit sind Frühdiagnose und Verlauf des Parkinson-Syndroms, sowie zerebellärer und spastischer Symptome möglich. Durch die Anwendung der Testbatterie für die kognitiven Funktionen können Frühdiagnose und Verlaufs kontrollen dementieller Syndrome erfolgen. Die Kognitotherapie findet bereits bei Frontalhirnverletzungen Anwendung.

## Kinematic analysis in evaluation of movement disorders

M. Berger, S. Mescheriakov, F. Gerstenbrand, I. Kozlovskaya\*, B. Babaev\*, A. Sokolov\* (Universitätsklinik für Neurologie, Innsbruck, and \*Institute of Biomedical Problems, Moscow)

Complex disturbances of coordination of aimed voluntary movements in microgravity, which are known as a hypogravitational ataxia (I. Kozlovskaya, 1983) and manifested by reduced accuracy and altered kinematics of aimed movements, are suggested to appear due to increased excitability of the motor control structures including increased activation of antagonists (I. Kozlovskaya 1983, 1993).

We studied single-joint pointing arm movements performed in real and simulated microgravity under different visual control conditions in order to detect the influence of the changed gravity environment on the kinematics of single-joint movements. Results of investigations during five days dry immersion experiment (four subjects) and nine days spaceflight (one cosmonaut) are presented.

Recording of three-dimensional arm position was conducted using MÖNIMIR equipment installed on board of the space station MIR and using the terrestrial version of the same equipment in the dry immersion facility of the Institute of Space Neurology in Innsbruck.

For processing of the data a mathematical model of the coordination of simple preprogrammed arm movements were used (S. Mescherikov, 1994). The model is based on the assumption that time profiles of the angular acceleration produced separately by agonist and antagonist muscles could be approximately described using Gaussian distribution functions and was proposed in order to separately estimate the angular acceleration produced by agonist and antagonist muscles (or muscle groups) during simple preprogrammed arm movements (i.e. one acceleration and one deceleration phase).

The following parameters of a position-time, velocity-time, and acceleration-time profiles were analyzed: movement duration (MD), peak velocity (PV), time to peak velocity (AP), time from peak velocity to the end of the movement (DP), peak acceleration (PA), peak deceleration (PD), time from the movement onset to the peak acceleration (AT), time from the peak deceleration to the end of movement (DT), time between peak acceleration and peak deceleration (ST).

A significant increase of the MD and decrease of resultant PV of the movement was observed in inflight sessions during the short-term spaceflight. Analysis of the acceleration-time profile showed that increase of MD was caused mostly by prolongation of the ST and covaries with the decrease of PA and PD values.

In the simulated microgravity experiment the increase of MD after 24 hours of immersion was observed. This effect was more pronounced in trials with limited visual control (no visual feedback of the arm position) and covaries with prolongation of the DP. Comparison of phases of the acceleration-time profile revealed higher contribution of the ST in the increase of MD. The data presented show that the real and sim-



# Wiener Medizinische Wochenschrift

Supplement Nr. 110

## II. Europäisches Symposium Quantifizierung von rehabilitativen Trainingsprogrammen

Wien, 2. bis 5. Juni 1994  
Kongreßzentrum Hofburg  
(Abstracts)

## II. Europäisches Symposium Quantifizierung von rehabilitativen Trainingsprogrammen. Wien, 2 bis 5. Juni 1994.

Ausgabe/Medienart  Artikel : Deutsch

Quelle: Wiener medizinische Wochenschrift 1994; 144 Suppl 110: 1-28

### II. Europäisches Symposium Quantifizierung von rehabilitativen Trainingsprogrammen : Wien, 2. bis 5. Juni 1994, Kongresszentrum Hofburg : (Abstracts).

Verlag: Wien : Blackwell, 1994.

Serien: [Wiener medizinische Wochenschrift](#), 144. Jahrg., suppl. nr. 110.; [WMW](#), 144. Jahrg., suppl. nr. 110.

Ausgabe/Medienart  Gedrucktes Buch : Tagungsband : Deutsch

### Abstracts / Wien, 2. bis 5. Juni 1994, Kongreßzentrum Hofburg.

Autor: [Europäisches Symposium Quantifizierung von Rehabilitativen Trainingsprogrammen \(2, 1994, Wien\)](#)

Verlag: Wien : Blackwell-MZV, 1994.

Serien: [@Wiener medizinische Wochenschrift](#), 110.; [Wiener medizinische Wochenschrift / Supplement](#), 110.

Ausgabe/Medienart  Gedrucktes Buch : Tagungsband : Deutsch